Author Index

AGARWAL S. *see* KESTLER D.P.
AGEA E. *see* SPINOZZI F.
AHERN M. *see* BECKMAN I.
AHLSTEDT S. *see* DAHLMAN-HÖGLUND A.
AKIMOTO Y. *see* IMAI Y.
ALZANI R. *see* ZINETTI M.
AMIRAYAN N., FURRIE E., DELEUIL F., MELLOR A., LESERMAN L. & MACHY P. Influence of MHC class I molecules on T-cell proliferation induced by CD3 or Thy-1 stimulation, 71
AN S. *see* ZENG L.
ANDERSSON J. *see* SANDER B.
ANDERSSON U. *see* SANDER B.
ANDONEGUI G., TREVANI A.S., DRAN G. & GEFFNER J.R. Effect of nitric oxide donors on neutrophil responses induced by immune complexes, 646
ARAMAKI Y. *see* MURAI M.
ARÁNGUEZ I. *see* TORRES C.
AWINO E. *see* BEMBRIDGE G.P.
AZUMA T. *see* KANEKO Y.

BABIUK L.A. *see* BACA-ESTRADA M.E.
BACA-ESTRADA M.E., LIANG X., BABIUK L.A. & YOO D. Induction of mucosal immunity in cotton rats to haemagglutinin–esterase glycoprotein of bovine coronavirus by recombinant adenovirus, 134
BACH J.-F. *see* PATRY C.
BACON M. *see* BINNS R.M.
BALDWIN C.L. *see* OKRAGLY A.J.
BALK H.P. *see* HOEFAKKER S.
BALOGH P. & KUMÁNOVICS A. Tissue-associated phenotypic heterogeneity of peripheral B cells in mice, 560
BAMBERG T. *see* MCGEE D.W.
BARANYI L. *see* KANEKO Y.
BARBACANE R.C. *see* CONTI P.
BASKETTER D.A. *see* DEARMAN R.J.
BECKMAN I., SHEPHERD D., FIRGAIRA F. & AHERN M. Age-related defects in CD2 receptor-induced activation in human T-cell subsets, 533
BEESON D.M.W. *see* MULCAHY A.F.
BEFUS A.D. *see* BISSONNETTE E.Y.
BELIA S. *see* SPINOZZI F.
BELLABY T. *see* ROBINSON K.
BEMBRIDGE G.P., MACHUGH N.D., MCKEEVER D., AWINO E., SOPP P., COLLINS R.A., GELDER K.I. & HOWARD C.J. CD45RO expression on bovine T cells: relation to biological function, 537
BERTOTTO A. *see* SPINOZZI F.
BHEEKHA ESCURA R., WASSERBAUER E., HAMMERSCHMID F., PEARCE A., KIDD P. & MUDDE G.C. Regulation and targeting of T-cell immune responses by IgE and IgG antibodies, 343
BIANCHI A.T.J. *see* KIMMAN T.G.
BIANCO N.E. *see* DE SANCTIS J.B.
BIBERFELD P. *see* CASTAÑOS-VELEZ E.
BILYK N. & HOLT P.G. Cytokine modulation of the immunosuppressive phenotype of pulmonary alveolar macrophage populations, 231
BINNS R.M., LICENCE S.T., WHYTE A., WILBY M., ROTHKÖTTER H.-J. & BACON M. Genetically determined CD45 variant of value in leucocyte tracing *in vivo* in the pig, 25
BISSONNETTE E.Y., CHIN B. & BEFUS A.D. Interferons differentially regulate histamine and TNF-α in rat intestinal mucosal mast cells, 12
BLANCA I. *see* DE SANCTIS J.B.
BODMER M.W. *see* MORGAN A.
BOERSMA W.J.A. *see* HOEFAKKER S.
BONATO V.L.D. *see* SILVA C.L.
BORREBAECK C.A.K. *see* MÅRTENSSON C.
BOUCHER W. *see* CONTI P.
BRAND D.D. *see* MIYAHARA H.
BRONDZ B.D., KAZANSKY D.B., CHERNYSHOVA A.D. & IVANOV V.S. Peptides of a major histocompatibility complex class I (K^b) molecule cause prolongation of skin graft survival and induce specific down-regulatory T cells demonstrable in the mixed lymphocyte reaction, 219
BRUNEAU G. *see* CONTI P.
BRYAN-POOLE N. *see* ZUCKERMAN S.H.

CALDER V.L. *see* GREENWOOD J.
CARLSSON R. *see* MÅRTENSSON C.
CASTAÑOS-VELEZ E., BIBERFELD P. & PATARROYO M. Extracellular matrix proteins and integrin receptors in reactive and non-reactive lymph nodes, 270
CATTANEO A. *see* MOZZANICA N.
CAULFIELD M.J. & STANKO D. T-cell dependent response to immune complexes abrogates B-cell unresponsiveness to pneumococcal cell wall polysaccharide, 331
CHADEE K. *see* SEGUIN R.
CHALLACOMBE S.J. *see* SWEET S.P.
CHAN W.C. *see* ROBINSON K.
CHAPLIN L. *see* MORGAN A.
CHEN M. *see* MULLER D.
see also TALMOR M.
CHEN X.-J., WIEDERMANN U., DAHLGREN U., HANSON L.Å & ENERBÄCK L. T-cell-independent and T-cell-dependent IgE responses to the nematode *Nippostrongylus brasiliensis*: comparison of serum IgE and mast-cell-bound IgE, 351
CHERNYSHOVA A.D. *see* BRONDZ B.D.
CHIN B. *see* BISSONNETTE E.Y.
CIPOLLARO DE L'ERO G. *see* GALDIERO M.
CLAASSEN E. *see* HOEFAKKER S.
CLERICI M. *see* MOZZANICA N.
CODNER M.A. *see* TALMOR M.
COE J.E., SCHELL R.F. & ROSS M.J. Immune response in the hamster: definition of a novel IgG not expressed in all hamster strains, 141
COLEMAN J.W. *see* TAYLOR A.M.
see also WILLIAMS C.M.M.
COLLINS R.A. *see* BEMBRIDGE G.P.
COLOVAI A.I. *see* HARRIS P.E.
COLSTON M.J. *see* MISRA N.
CONTI P., BOUCHER W., LETOURNEAU R., FELICIANI C., REALE M., BARBACANE R.C., VLAGOPOULOS P., BRUNEAU G., THIBAULT J. & THEOHARIDES T.C. Monocyte chemotactic protein-1 provokes mast cell aggregation and [^{3}H]5HT release, 434
COZZI E. *see* ZINETTI M.
CUMBERBATCH M. *see* HOPE J.C.
CUNNINGHAM A.C. & KIRBY J.A. Regulation and function of adhesion molecule expression by human alveolar epithelial cells, 279
CUNNINGHAM A.C. *see* WILSON J.L.

DA SILVA M.F. *see* SILVA C.L.
DAHLGREN U. *see* CHEN X.-J.
see also DAHLMAN-HÖGLUND A.
DAHLMAN-HÖGLUND A., DAHLGREN U., AHLSTEDT S., HANSON L.Å & TELEMO E. Bystander suppression of the immune response to human serum albumin in rats fed ovalbumin, 128

DAI R. & STREILEIN J.W. Sensitizing capacity of Langerhans' cells obtained from ultraviolet-B-exposed murine skin, 661
DAMM O. *see* SANDER B.
DÄUBENER W., MACKENZIE C. & HADDING U. Establishment of T-helper type 1- and T-helper type 2-like human *Toxoplasma* antigen-specific T-cell clones, 79
DAVIES M.E. *see* HORNER A.
DAY N.K. *see* KIZAKI T.
DE BRUIN T.M.G. *see* KIMMAN T.G.
DE SANCTIS J.B., BLANCA I. & BIANCO N.E. Expression of different lipoprotein receptors in natural killer cells and their effect on natural killer proliferative and cytoxic activity, 399
DEARMAN R.J. *see* HOPE J.C.
DEARMAN R.J., BASKETTER D.A. & KIMBER I. Differential cytokine production following chronic exposure of mice to chemical respiratory and contact allergens, 545
DELEUIL F. *see* AMIRAYAN N.
DEMAISON C., RAVIRAJAN C.T., ISENBERG D.A. & ZOUALI M. Analysis of variable region genes encoding anti-Sm and anti-cardiolipin antibodies from a systemic lupus erythematosus patient, 487
DEMITRI M.T. *see* ZINETTI M.
DENSEN P. *see* MAVES K.K.
DIAMOND A.G. *see* MULCAHY A.F.
DIAZ-MITOMA F. *see* KRYWORUCKHO M.
DONNARUMMA G. *see* GALDIERO M.
DRAN G. *see* ANDONEGUI G.
DYBWAD A. *see* ØSTENSTAD B.

EMOTO M. *see* KITA E.
EMTAGE J.S. *see* MORGAN A.
ENERBÄCK L. *see* CHEN X.-J.
ERIKSSON H. & SJÖGREN H.O. Inhibition of activation of human T lymphocytes by the complement C1 esterase inhibitor, 304
EVANS G.F. *see* ZUCKERMAN S.H.

FANTUZZI G. *see* ZINETTI M.
FELICIANI C. *see* CONTI P.
FERRARIO E. *see* MOZZANICA N.
FIELDING I. *see* HOPE J.C.
FINZI A.F. *see* MOZZANICA N.
FIRGAIRA F. *see* BECKMAN I.
FØRRE Ø. *see* ØSTENSTAD B.
FRANZ B. *see* HORNER A.
FRATELLI M. *see* ZINETTI M.
FURRIE E. *see* AMIRAYAN N.
FURRIE E., TURNER M.W. & STROBEL S. Partial characterization of a circulating tolerogenic moiety which, after a feed of ovalbumin, suppresses delayed-type hypersensitivity in recipient mice, 480

GALDIERO F. *see* GALDIERO M.
GALDIERO M., CIPOLLARO DE L'ERO G., DONNARUMMA G., MARCATILI A. & GALDIERO F. Interleukin-1 and interleukin-6 gene expression in human monocytes stimulated with *Salmonella tymphimurium* porins, 612
GALLI G. *see* ZINETTI M.
GALLI S.J. *see* TAYLOR A.M.
GAVIN A.L., SNIDER J., HULETT M.D., MCKENZIE I.F.C. & HOGARTH P.M. Expression of recombinant soluble FcεRI: function and tissue distribution studies, 392
GEFFNER J.R. *see* ANDONEGUI G.
GELDER K.I. *see* BEMBRIDGE G.P.
GERSHON D. *see* ROBY K.F.
GHEZZI P. *see* ZINETTI M.
GLASEBROOK A.L. *see* ZUCKERMAN S.H.
GOETZL E.J. *see* ZENG L.
GOLDBERG I. *see* OREN R.
GOOD R.A. *see* KIZAKI T.
GREENWOOD J., WANG Y. & CALDER V.L. Lymphocyte adhesion and transendothelial migration in the central nervous system: the role of LFA-1, ICAM-1, VLA-4 and VCAM-1, 408
GRIGNANI F. *see* SPINOZZI F.
GUENTHNER S.T. *see* MAVES K.K.
GUILLET J.-G. *see* PLASMAN N.

HADDING U. *see* DÄUBENER W.
HÅKANSSON L. *see* SANDER B.
HALL R.E. *see* KESTLER D.P.
HÄLLSTRÖM O. *see* KERTTULA T.O.
HAMMERSCHMID F. *see* BHEEKHA ESCURA R.
HANBY-FLARIDA M. *see* OKRAGLY A.J.
HANSON L.Å *see* CHEN X.-J.
see also DAHLMAN-HÖGLUND A.
HARPER A.D. *see* TALMOR M.
HARRIS C.L. & MORGAN B.P. Characterization of a glycosyl-phosphatidylinositol anchor-deficient subline of Raji cells. An analysis of the functional importance of complement inhibitors on the Raji cell line, 311
HARRIS D.P. *see* ROMÁN E.
HARRIS P.E., COLOVAI A.I., MAFFEI A., LIU Z. & SUCIU FOCA N. Major histocompatibility complex class I presentation of exogenous and endogenous protein-derived peptides by a transfected human monocyte cell line, 606
HEEG K. *see* MARTINEZ-KINADER B.
HERBELIN A. *see* PATRY C.
HERNÁNDEZ-PANDO R., OROZCO H. & MANCILLA R. T-cell lung granulomas induced by sepharose-coupled *Mycobacterium tuberculosis* protein antigens: immunosuppressive phenomena reversed with cyclophosphamide and indomethacin, 506
HILDEMAN D. *see* MULLER D.
HIRANO H. *see* IMAI Y.
HIROKAWA M., KITABAYASHI A., KUROKI J. & MIURA A.B. Signal transduction by B7/BB1 expressed on activated T lymphocytes: cross-linking of B7/BB1 induces protein tyrosine phosphorylation and synergizes with signalling through T-cell receptor/CD3, 155
HIROMATSU K. *see* USAMI J.
HOEFAKKER S., BALK H.P., BOERSMA W.J.A., VAN JOOST TH., NOTTEN W.R.F. & CLAASSEN E. Migration of human antigen-presenting cells in a human skin graft onto nude mice model after contact sensitization, 296
HOFFMAN L.A. *see* TALMOR M.
HOGARTH P.M. *see* GAVIN A.L.
HOLT P.G. *see* BILYK N.
HOLTER W. *see* SCHWARZ M.
HOPE J.C., CUMBERBATCH M., FIELDING I., DEARMAN R.J., KIMBER I. & HOPKINS S.J. Identification of dendritic cells as a major source of interleukin-6 in draining lymph nodes following skin sensitization of mice, 441
HOPKINS S.J. *see* HOPE J.C.
HORNER A., DAVIES M.E. & FRANZ B. Chrondocyte–peripheral blood mononuclear cell interactions: the role of ICAM-1, 584
HOWARD C.J. *see* BEMBRIDGE G.P.
HOWIE D. *see* SANDILANDS G.P.
HULETT M.D. *see* GAVIN A.L.
HUNT J.S. *see* ROBY K.F.

IFVERSEN P. *see* MÅRTENSSON C.
IMAI Y., AKIMOTO Y., MIZUOCHI S., KIMURA T., HIRANO H. & IRIMURA T. Restricted expression of galactose/*N*-acetylgalactosamine-specific macrophage C-type lectin to connective tissue and to metastatic lesions in mouse lung, 591
INAGAKI H. *see* USAMI J.
IRIMURA T. *see* IMAI Y.
ISENBERG D.A. *see* DEMAISON C.
IVANOV V.S. *see* BRONDZ B.D.
IVANYI J. *see* ROMÁN E.

JANUARIO A. *see* SILVA C.L.
JONES J. & MORGAN B.P. Apoptosis is associated with reduced expression of complement regulatory molecules, adhesion molecules and other receptors on polymorphonuclear leucocytes: functional relevance and role in inflammation, 651
JONES N.D. *see* MORGAN A.
JURCEVIC S. *see* ROMÁN E.

KANAMORI H., KODAMA T. & YAZAKI Y. The 5′ coding sequence of IL-2 receptor α chain mRNA mediates mRNA stabilization by HTLV-1 Rex, 551
KANEKO Y., OKADA N., BARANYI L., AZUMA T. & OKADA H. Antagonistic peptides against human anaphylatoxin C5a, 149
KANG A.H. *see* MIYAHARA H.
KAPP J.A. *see* ZIMECKI M.
KASHIBA S. *see* KITA E.
KAZANSKY D.B. *see* BRONDZ B.D.
KELLER K. *see* SEGUIN R.
KERTTULA T.O., HÄLLSTRÖM O. & MÄKI M. Phenotypical characterization of peripheral blood T cells in patients with coeliac disease: elevation of antigen-primed $CD45RO^+$ T lymphocytes, 104
KESTLER D.P., AGARWAL S. & HALL R.E. Up-regulation of cytokine mRNA in human monocytes and myeloid cell lines by the differentiation/activation factor p48, 463
KIDD P. *see* BHEEKHA ESCURA R.
KILSHAW P.J. *see* PENNEY L.
KIMBER I. *see* DEARMAN R.J.
see also HOPE J.C.
KIMMAN T.G., DE BRUIN T.M.G., VOERMANS J.J.M., PEETERS B.P.H. & BIANCHI A.T.J. Development and antigen specificity of the lymphoproliferation response of pigs to pseudorabies virus: dichotomy between secondary B- and T-cell responses, 372
KIMURA T. *see* IMAI Y.
KIRBY J.A. *see* CUNNINGHAM A.C.
KIRBY J.A. *see* WILSON J.L.
KITA E., EMOTO M., NISHIKAWA F., YOSHIKAI Y. & KASHIBA S. Conversion of *Salmonella typhimurium* to L-forms contributes to the maintenance of acquired immunity against murine typhoid, 206
KITABAYASHI A. *see* HIROKAWA M.
KIZAKI T., YAMASHITA H., OH-ISHI S., DAY N.K., GOOD R.A. & OHNO H. Immunomodulation by cells of mononuclear phagocyte lineage in acute cold-stressed or cold-acclimatized mice, 456
KNAPP W. *see* SCHWARZ M.
KODAMA T. *see* KANAMORI H.
KOIDE Y., YOSHIDA A., UCHIJIMA M. & YOSHIDA T.O. Unimpaired clearance of *Mycobacterium bovis* BCG infection in selectively T-cell anergic TCR-Vβ8.2 transgenic mice, 499
KONG Y.-Y. *see* MURAKAMI Y.
KONING F. *see* ORSINI D.L.M.
KOOY Y.M.C. *see* ORSINI D.L.M.
KOPOLOVIC J. *see* OREN R.
KRYWORUCKHO M., DIAZ-MITOMA F. & KUMAR A. CD44 isoforms containing exons V6 and V7 are differentially expressed on mitogenically stimulated normal and Epstein–Barr virus-transformed human B cells, 41
KUMÁNOVICS A. *see* BALOGH P.
KUMAR A. *see* KRYWORUCKHO M.
KUROKI J. *see* HIROKAWA M.

LAMB J.R. *see* O'BRIEN R.M.
LANGERMANS J.A.M. *see* SAMSOM J.N.
LAUFER J. *see* OREN R.
LEA T. *see* ØSTENSTAD B.
LEHUEN A. *see* PATRY C.
LESERMAN L. *see* AMIRAYAN N.
LETOURNEAU R. *see* CONTI P.
LIANG X. *see* BACA-ESTRADA M.E.
LICENCE S.T. *see* BINNS R.M.
LIMA V.M.F. *see* SILVA C.L.
LIPFORD G.B. *see* MARTINEZ-KINADER B.
LIU Z. *see* HARRIS P.E.
LOWRIE D.B. *see* SILVA C.L.
LU P. & SHAROM F.J. Gangliosides are potent immunosuppressors of IL-2-mediated T-cell proliferation in a low protein environment, 356
LYCKE N. *see* VAJDY M.

MA D. & NIEDERKORN J.Y. Transforming growth factor-β down-regulates major histocompatibility complex class I antigen expression and increases the susceptibility of uveal melanoma cells to natural killer cell-mediated cytolysis, 263
MACDONALD T.T. *see* PENNEY L.
MACHUGH N.D. *see* BEMBRIDGE G.P.
MACHY P. *see* AMIRAYAN N.
MACKENZIE C. *see* DÄUBENER W.
MACSWEEN R.N.M. *see* SANDILANDS G.P.
MAEDA K. *see* USAMI J.
MAEDA T. *see* MURAKAMI Y.
MAFFEI A. *see* HARRIS P.E.
MAJDIC O. *see* SCHWARZ M.
MÄKI M. *see* KERTTULA T.O.
MANCILLA R. *see* HERNÁNDEZ-PANDO R.
MARCATILI A. *see* GALDIERO M.
MÅRTENSSON C., IFVERSEN P., BORREBAECK C.A.K. & CARLSSON R. Enhancement of specific immunoglobulin production in SCID-hu-PBL mice after *in vitro* priming of human B cells with superantigen, 224
MARTIN S.J. *see* VYAKARNAM A.
MARTINEZ-KINADER B., LIPFORD G.B., WAGNER H. & HEEG K. Sensitization of MHC class I-restricted T cells to exogenous proteins: evidence for an alternative class I-restricted antigen presentation pathway, 287
MATEAR P.M. *see* VYAKARNAM A.
MATSUMOTO Y. *see* USAMI J.
MATSUZAKI G. *see* SONODA K.
MAVES K.K., GUENTHNER S.T., DENSEN P., MOSER D.R. & WEILER J.M. Cloning and characterization of the cDNA encoding guinea-pig properdin: a comparison of properdin from three species, 475
MAZURIER J. *see* ZIMECKI M.
MCGEE D.W., BAMBERG T., VITKUS S.J.D. & MCGHEE J.R. A synergistic relationship between TNF-α, IL-1β and TGF-β1 on IL-6 secretion by the IEC-6 intestinal epithelial cell line, 6
MCGHEE J.R. *see* MCGEE D.W.
MCKEEVER D. *see* BEMBRIDGE G.P.
MCKENZIE I.F.C. *see* GAVIN A.L.
MCLAREN A.P. *see* SANDILANDS G.P.
MEGYERI P., PABST K.M. & PABST M.J. Serine protease inhibitors block priming of monocytes for enhanced release of superoxide, 629
MELLOR A. *see* AMIRAYAN N.
MIGLIORATI G. *see* SPINOZZI F.
MINTO M. *see* ZINETTI M.
MISRA N., MURTAZA A., WALKER B., NARAYAN N.P.S., MISRA R.S., RAMESH V., SINGH S., COLSTON M.J. & NATH I. Cytokine profile of circulating T cells of leprosy patients reflects both indiscriminate and polarized T-helper subsets: T-helper phenotype is stable and uninfluenced by related antigens of *Mycobacteria leprae*, 97
MISRA R.S. *see* MISRA N.
MIURA A.B. *see* HIROKAWA M.
MIYAHARA H., MYERS L.K., ROSLONIEC E.F., BRAND D.D., SEYER J.M., STUART J.M. & KANG A.H. Identification and characterization of a major tolerogenic T-cell epitope of type II collagen that suppresses arthritis in B10.RIII mice, 110
MIZUOCHI S. *see* IMAI Y.
MONTEIRO R.C. *see* PATRY C.
MORACA R. *see* SPINOZZI F.
MORENO C. *see* ROMÁN E.
MORGAN A., JONES N.D., NESBITT A.M., CHAPLIN L., BODMER M.W. & EMTAGE J.S. The N-terminal end of the C_H2 domain of chimeric human IgG1 anti-HLA-DR is necessary for C1q, FcγRI and FcγRII binding, 319
MORGAN B.P. *see* HARRIS C.L.
see also JONES J.
MOSER D.R. *see* MAVES K.K.
MOZZANICA N., CATTANEO A., TRABATTONI D., FINZI A.F., SCHMITT E., FERRARIO E., CLERICI M., VIGNATI G. & VILLA M.L. Production of type-1 and type-2 cytokines by peripheral blood mononuclear cells of psoriatic patients, 422
MUDDE G.C. *see* BHEEKHA ESCURA R.
MUKASA A. *see* SONODA K.

MULCAHY A.F., BEESON D.M.W., WILLCOX N. & DIAMOND A.G. Presentation of endogenous acetylcholine receptor antigen to a specific $CD4^+$ T-cell line by a transfected B-cell line, 116
MULLER D., CHEN M., VIKINGSSON A., HILDEMAN D. & PEDERSON K. Oestrogen influences $CD4^+$ T-lymphocyte activity *in vivo* and *in vitro* in β_2-microglobulin-deficient mice, 162
MURAI M., ARAMAKI Y. & TSUCHIYA S. Identification of the serum factor required for liposome-primed activation of mouse peritoneal macrophages. Modified α_2-macroglobulin enhances Fcγ receptor-mediated phagocytosis of opsonized sheep red blood cells, 64
MURAI M., ARAMAKI Y. & TSUCHIYA S. Modification of α_2-macroglobulin into a macrophage-activating factor through the action of liposome-stimulated B-cell membranous glycosidases, 58
MURAKAMI Y., KONG Y.-Y., NISHIMURA Y., NOMOTO K., UMESUE M., OMOTO K., MAEDA T. & NOMOTO K. Prevention of anti-T-cell receptor $\alpha\beta$ monoclonal antibody-induced side-effects by treatment with cyclosporin A without interference of monoclonal antibody-induced immunosuppression in mice, 238
MURTAZA A. *see* MISRA N.
MYERS L.K. *see* MIYAHARA H.

NARAYAN N.P.S. *see* MISRA N.
NATH I. *see* MISRA N.
NESBITT A.M. *see* MORGAN A.
NICHOLSON I. *see* O'BRIEN R.M.
NICOLETTI I. *see* SPINOZZI F.
NIEDERKORN J.Y. *see* MA D.
NISHIKAWA F. *see* KITA E.
NISHIMURA Y. *see* MURAKAMI Y.
NOMOTO K. *see* MURAKAMI Y.
see also SONODA K.
NOTTEN W.R.F. *see* HOEFAKKER S.

O'BRIEN R.M., THOMAS W.R., NICHOLSON I., LAMB J.R. & TAIT B.D. An immunogenetic analysis of the T-cell recognition of the major house dust mite allergen Der p 2: identification of high- and low-responder HLA-DQ alleles and localization of T-cell epitope, 176
OH-ISHI S. *see* KIZAKI T.
OHNO H. *see* KIZAKI T.
OKADA H. *see* KANEKO Y.
OKADA N. *see* KANEKO Y.
OKRAGLY A.J., HANBY-FLARIDA M. & BALDWIN C.L. Monocytes control $\gamma\delta$ T-cell responses by a secreted product, 599
OMOTO K. *see* MURAKAMI Y.
OREN R., LAUFER J., GOLDBERG I., KOPOLOVIC J., WALDHERR R. & PASSWELL J.H. C3, C4, factor B and HLA-DRα mRNA expression in renal biopsy specimens from patients with IgA nephropathy, 575
OROZCO H. *see* HERNÁNDEZ-PANDO R.
ORSINI D.L.M., KOOY Y.M.C., VAN DER TOL M.A., STRUYK L., VAN DEN ELSEN P. & KONING F. T-cell receptor usage of interleukin-2-responsive peripheral $\gamma\delta$ T cells, 385
ØSTENSTAD B., DYBWAD A., LEA T., FØRRE Ø., VINJE O. & SIOUD M. Evidence for monoclonal expansion of synovial T cells bearing Vα2.1/Vβ5.5 gene segments and recognizing a synthetic peptide that shares homology with a number of putative autoantigens, 168

PABST K.M. *see* MEGYERI P.
PABST M.J. *see* MEGYERI P.
PASSWELL J.H. *see* OREN R.
PATARROYO M. *see* CASTAÑOS-VELEZ E.
PATRY C., HERBELIN A., LEHUEN A., BACH J.-F. & MONTEIRO R.C. Fcα receptors mediate release of tumour necrosis factor-α and interleukin-6 by human monocytes following receptor aggregation, 1
PEARCE A. *see* BHEEKHA ESCURA R.
PEDERSON K. *see* MULLER D.
PEETERS B.P.H. *see* KIMMAN T.G.
PENNEY L., KILSHAW P.J. & MACDONALD T.T. Regional variation in the proliferative rate and lifespan of $\alpha\beta$ TCR^+ and $\gamma\delta$ TCR^+ intraepithelial lymphocytes in the murine small intestine, 212
PIETERS R.H.H. *see* SCHIELEN P.
PIETRO R.L.R. *see* SILVA C.L.
PLASMAN N., GUILLET J.-G. & VRAY B. Impaired protein catabolism in *Trypanosoma cruzi*-infected macrophages: possible involvement in antigen presentation, 636

RAHMAN D. *see* SWEET S.P.
RAMESH V. *see* MISRA N.
RAVIRAJAN C.T. *see* DEMAISON C.
REALE M. *see* CONTI P.
RICCARDI C. *see* SPINOZZI F.
ROBINSON K., BELLABY T., CHAN W.C. & WAKELIN D. High levels of protection induced by a 40-mer synthetic peptide vaccine against the intestinal nematode parasite *Trichinella spiralis*, 495
ROBY K.F., YANG Y., GERSHON D. & HUNT J.S. Cellular distribution of proteasome subunit *Lmp7* mRNA and protein in human placentas, 469
ROMÁN E., HARRIS D.P., JURCEVIC S., IVANYI J. & MORENO C. H-2-associated effects of flanking residues on the recognition of a permissive mycobacterial T-cell epitope, 183
ROSLONIEC E.F. *see* MIYAHARA H.
ROSS M.J. *see* COE J.E.
ROTHKÖTTER H.-J. *see* BINNS R.M.
ROTTEVEEL F.T.M., VAN DOORNMALEN A.M. & VAN DUIN M. sLex is not responsible for the interaction of sLex-positive memory T lymphocytes with E-selectin 34
RUBIO N. *see* TORRES C.

SAMSOM J.N., LANGERMANS J.A.M., SAVELKOUL H.F.J. & VAN FURTH R. Tumour necrosis factor, but not interferon-γ, is essential for acquired resistance to *Listeria monocytogenes* during a secondary infection in mice, 256
SANDER B., SKANSÉN-SAPHIR U., DAMM O., HÅKANSSON L., ANDERSSON J. & ANDERSSON U. Sequential production of Th1 and Th2 cytokines in response to live bacillus Calmette–Guérin, 512
SANDILANDS G.P., MCLAREN A.P., HOWIE D. & MACSWEEN R.N.M. Occult expression of CD32 (FcγRII) in normal human peripheral blood mononuclear cells, 525
SAVELKOUL H.F.J. *see* SAMSOM J.N.
SCHELL R.F. *see* COE J.E.
SCHIELEN P., VAN RODIJNEN W., PIETERS R.H.H. & SEINEN W. Hexachlorobenzene treatment increases the number of splenic B-1-like cells and serum autoantibody levels in the rat, 568
SCHMITT E. *see* MOZZANICA N.
SCHWARZ M., MAJDIC O., KNAPP W. & HOLTER W. High-level IL-10 production by monoclonal antibody-stimulated human T cells, 364
SEGUIN R., KELLER K. & CHADEE K. *Entamoeba histolytica* stimulates the unstable transcription of c-fos and tumour necrosis factor-α mRNA by protein kinase C signal transduction in macrophages, 49
SEINEN W. *see* SCHIELEN P.
SEYER J.M. *see* MIYAHARA H.
SHAROM F.J. *see* LU P.
SHEPHERD D. *see* BECKMAN I.
SHORT L. *see* ZUCKERMAN S.H.
SILVA C.L., PIETRO R.L.R., JANUARIO A., BONATO V.L.D., LIMA V.M.F., DA SILVA M.F. & LOWRIE D.B. Protection against tuberculosis by bone marrow cells expressing mycobacterial hsp65, 519
SINGH S. *see* MISRA N.
SIOUD M. *see* ØSTENSTAD B.
SJÖGREN H.O. *see* ERIKSSON H.
SKANSÉN-SAPHIR U. *see* SANDER B.
SNIDER J. *see* GAVIN A.L.
SONODA K., MUKASA A., MATSUZAKI G. & NOMOTO K. The induction of renal autoantigen-specific T cells by a local *Listeria monocytogenes* infection, 190

SOPP P. *see* BEMBRIDGE G.P.
SPIK G. *see* ZIMECKI M.
SPINOZZI F., NICOLETTI I., AGEA E., BELIA S., MORACA R., MIGLIORATI G., RICCARDI C., GRIGNANI F. & BERTOTTO A. IL-4 is able to reverse the CD2-mediated negative apoptotic signal to $CD4^-CD8^+$ $\alpha\beta$ and/or $\gamma\delta$ T lymphocytes, 379
STANKO D. *see* CAULFIELD M.J.
STEINMAN R.M. *see* TALMOR M.
STREILEIN J.W. *see* DAI R.
STROBEL S. *see* FURRIE E.
STRUYK L. *see* ORSINI D.L.M.
STUART J.M. *see* MIYAHARA H.
SUCIU FOCA N. *see* HARRIS P.E.
SUZUKI T. *see* USAMI J.
SWEET S.P., RAHMAN D. & CHALLACOMBE S.J. IgA subclasses in HIV disease: dichotomy between raised levels in serum and decreased secretion rates in saliva, 556

TAIT B.D. *see* O'BRIEN R.M.
TALMOR M., STEINMAN R.M., CODNER M.A., CHEN M., HARPER A.D., WITMER PACK M.D. & HOFFMAN L.A. Bone marrow-derived chimerism in non-irradiated, cyclosporin-treated rats receiving microvascularized limb transplants: evidence for donor-derived dendritic cells in recipient lymphoid tissues, 448
TAYLOR A.M., GALLI S.J. & COLEMAN J.W. Stem-cell factor, the kit ligand, induces direct degranulation of rat peritoneal mast cells *in vitro* and *in vivo*: dependence of the *in vitro* effect on period of culture and comparisons of stem-cell factor with other mast cell-activating agents, 427
TELEMO E. *see* DAHLMAN-HÖGLUND A.
THEOHARIDES T.C. *see* CONTI P.
THIBAULT J. *see* CONTI P.
THOMAS W.R. *see* O'BRIEN R.M.
TORRES C., ARÁNGUEZ I. & RUBIO N. Expression of interferon-γ receptors on murine oligodendrocytes and its regulation by cytokines and mitogens, 250
TRABATTONI D. *see* MOZZANICA N.
TREVANI A.S. *see* ANDONEGUI G.
TSUCHIYA S. *see* MURAI M.
TURNER M.W. *see* FURRIE E.

UCHIJIMA M. *see* KOIDE Y.
UMESUE M. *see* MURAKAMI Y.
USAMI J., HIROMATSU K., MATSUMOTO Y., MAEDA K., INAGAKI H., SUZUKI T. & YOSHIKAI Y. A protective role of $\gamma\delta$ T cells in primary infection with *Listeria monocytogenes* in autoimmune non-obese diabetic mice, 199

VAJDY M. & LYCKE N. Mucosal memory B cells retain the ability to produce IgM antibodies 2 years after oral immunization, 336
VAN DEN ELSEN P. *see* ORSINI D.L.M.
VAN DER TOL M.A. *see* ORSINI D.L.M.
VAN DOORNMALEN A.M. *see* ROTTEVEEL F.T.M.
VAN DUIN M. *see* ROTTEVEEL F.T.M.
VAN FURTH R. *see* SAMSOM J.N.
VAN JOOST TH. *see* HOEFAKKER S.
VAN RODIJNEN W. *see* SCHIELEN P.
VIGNATI G. *see* MOZZANICA N.
VIKINGSSON A. *see* MULLER D.
VILLA M.L. *see* MOZZANICA N.
VINJE O. *see* ØSTENSTAD B.
VITKUS S.J.D. *see* MCGEE D.W.
VLAGOPOULOS P. *see* CONTI P.
VOERMANS J.J.M. *see* KIMMAN T.G.
VRAY B. *see* PLASMAN N.
VYAKARNAM A., MATEAR P.M., MARTIN S.J. & WAGSTAFF M. Th1 cells specific for HIV-1 gag p24 are less efficient than Th0 cells in supporting HIV replication, and inhibit virus replication in Th0 cells, 85

WAGNER H. *see* MARTINEZ-KINADER B.
WAGSTAFF M. *see* VYAKARNAM A.
WAKELIN D. *see* ROBINSON K.
WALDHERR R. *see* OREN R.
WALKER B. *see* MISRA N.
WANG Y. *see* GREENWOOD J.
WASSERBAUER E. *see* BHEEKHA ESCURA R.
WEILER J.M. *see* MAVES K.K.
WHYTE A. *see* BINNS R.M.
WIEDERMANN U. *see* CHEN X.-J.
WILBY M. *see* BINNS R.M.
WILLCOX N. *see* MULCAHY A.F.
WILLIAMS C.M.M. & COLEMAN J.W. Induced expression of mRNA for IL-5, IL-6, TNF-α, MIP-2 and IFN-γ in immunologically activated rat peritoneal mast cells: inhibition by dexamethasone and cyclosporin A, 244
WILSON J.L., CUNNINGHAM A.C. & KIRBY J.A. Alloantigen presentation by B cells: analysis of the requirement for B-cell activation, 325
WITMER-PACK M.D. *see* TALMOR M.

YAMASHITA H. *see* KIZAKI T.
YANG Y. *see* ROBY K.F.
YAZAKI Y. *see* KANAMORI H.
YOO D. *see* BACA-ESTRADA M.E.
YOSHIDA A. *see* KOIDE Y.
YOSHIDA T.O. *see* KOIDE Y.
YOSHIKAI Y. *see* KITA E.
YOSHIKAI Y. *see* USAMI J.

ZENG L., AN S. & GOETZL E.J. Independent down-regulation of EP_2 and EP_3 subtypes of the prostaglandin E_2 receptors on U937 human monocytic cells, 620
ZIMECKI M., MAZURIER J., SPIK G. & KAPP J.A. Human lactoferrin induces phenotypic and functional changes in murine splenic B cells, 122
ZINETTI M., GALLI G., DEMITRI M.T., FANTUZZI G., MINTO M., GHEZZI P., ALZANI R., COZZI E. & FRATELLI M. Chlorpromazine inhibits tumour necrosis factor synthesis and cytotoxicity *in vitro*, 416
ZOUALI M. *see* DEMAISON C.
ZUCKERMAN S.H., BRYAN-POOLE N., EVANS G.F., SHORT L. & GLASEBROOK A.L. *In vivo* modulation of murine serum tumour necrosis factor and interleukin-6 levels during endotoxemia by oestrogen agonists and antagonists, 18

Subject Index

Acetylcholine receptor antigen:
presentation to T-cell line by a transfected B-cell line, 116
Adhesion molecules:
expression by human alveolar epithelial cells, 279
Alloantigen presentation by B cells, 325
Anaphylatoxin C5a, human:
antagonistic peptides, 149
Antigen presentation:
in *Trypanosoma cruzi*-infected macrophages, 636
Apoptosis, polymorphonuclear leucocyte:
and receptor loss, 651

B-cell activation:
by hexachlorobenzene, 568
B-cell heterogeneity in mice, 560
B-memory cells, mucosal:
IgM antibody production, 336
Bacillus Calmette–Guérin infection:
in anergic transgenic mice, 499
Bone marrow chimerism:
in transplanted rats, 448
Bystander suppression of immune responses, 128

C1 esterase inhibitor:
inhibition of T-cell activation, 304
CD44 isoform:
in normal and transformed human B cells, 41
CD45RO expression:
on bovine T cells, 537
Class I-restricted presentation of exogenous proteins, 287
Contact sensitization:
on human xenografted skin, 296
Cyclosporin A:
prevention of anti-T-cell-receptor $\alpha\beta$ side-effects, 238
Cytokine mRNA:
up-regulation by differentiation/activation factor p48, 463
Cytokine production in mast cells:
inhibition by dexamethasone and cyclosporin A, 244
Cytokine production:
bacillus Calmette–Guérin induced, 512
following exposure to chemical allergens, 545
Cytokines:
modulation of alveolar macrophage immunosuppression, 231
production by T cells from patients with psoriasis, 422

Delayed-type hypersensitivity:
characterization of moiety of ovalbumin which suppresses, 480

Entamoeba histolytica:
stimulation of cytokine gene expression in macrophages, 49
E-selectin:
interaction with sLex memory T cells, 34
Extracellular matrix proteins and integrin receptors:
in lymph nodes, 270

Fcα receptors:
mediation of cytokine production, 1
FcεRI:
function and tissue distribution, 392
FcγRII expression, occult:
in normal blood mononuclear cells, 525

Gangliosides:
immunosuppression of interleukin-2-mediated T-cell proliferation, 356
Glycosyl phosphatidylinositol anchor-deficient Raji cells, 311

Human IgG1, chimeric:
C1q and FcγR binding sites, 319

ICAM-1:
role in chondrocyte–peripheral blood mononuclear cell interactions, 584
IgA nephropathy, C3, C4, factor B and HLA-DRα mRNA expression, 575
IgA subclasses:
in human immunodeficiency virus, 556
IgE responses to *Nippostongylus brasiliensis*:
T-cell dependent and independent, 351
IgG, novel:
hamster, 141
Immune complexes:
abrogation of B-cell unresponsiveness, 331
Immunomodulation:
induced by cold stress, 456
Interferon-γ receptors, on oligodendrocytes:
regulation by cytokines and mitogens, 250
Interferons:
differential regulation of histamine and tumour necrosis factor-α in mast cells, 12
Interleukin-1 and interleukin-6 expression in human monocytes stimulated with *Salmonella typhimurium* porins, 612
Interleukin-2 receptor α chain:
stabilization, 551
Interleukin-4:
reversal of CD2-mediated negative apoptotic signal, 379
Interleukin-6 secretion, by intestinal epithelial cell line:
synergistic relationship of cytokines on, 6
Interleukin-6:
production by dendritic cells, 441
Interleukin-10:
high-level production by antibody-stimulated human T cells, 364
Intraepithelial lymphocytes:
variation in proliferative rates and lifespan, 212

Lactoferrin, human:
effects on murine B cells, 122
Langerhans' cells, ultraviolet exposed:
sensitizing capacity, 661
Leucocyte tracing, pig:
using CD45 polymorphism, 25
Lipoprotein receptors and natural killer function, 399
Lymphocyte adhesion and migration, in CNS:
role of adhesion molecules, 408

α_2-macroglobulin:
enhancement of Fcγ receptor-mediated phagocytosis, 64
modification by glycosidases to a macrophage-activating factor, 58
Macrophage C-type lectin:
restricted expression, 591
Major histocompatibility complex class I (K^b) peptides:
effects on skin graft survival and mixed lymphocyte reaction, 219
Major histocompatibility complex class I antigen presentation:
by human monocytes, 606
Major histocompatibility complex class I molecules:
influence on T-cell proliferation, 71
Monocyte chemotactic protein-1:
promotion of histamine and 5HT-releasing factor for mast cells, 434

Monocyte priming:
blocking by protease inhibitors, 629
Mucosal immunity:
to bovine coronavirus, 134

Nitric oxide donors:
effects on neutrophil responses, 646

Oestrogen:
influence on $CD4^+$ T cells, 162

Properdin, pig:
cloning and characterization of cDNA, 475
Prostaglandin receptors:
down-regulation, 620
Proteasome subunit *Lmp7* mRNA and protein:
cellular distribution, 469
Pseudorabies virus:
immune responses in pigs, 372

Salmonella typhimurium, immunity to:
maintained by L forms of *Salmonella typhimurium*, 206
Signal transduction:
by B7/BB1 on activated T cells, 155
Stem-cell factor:
activation of mast cells, 427
Superantigen:
enhancement of immunoglobulin production in SCID mice following priming of B cells, 224
Synthetic peptide vaccine:
against *Trichinella spiralis*, 495

T cells, autoreactive:
induced by *Listeria monocytogenes* infection, 190
T cells, phenotypical characterization:
in patients with coeliac disease, 104
T cells, synovial fluid:
monoclonal expansion of, 168
γ/δ T cells:
T-cell receptor usage, 385
control of responses by monocytes, 599
in non-obese diabetic mice, 199
T-cell activation, CD2-induced:
in aged subjects, 533
T-cell clones, T-helper type 1- and T-helper type 2-like human:
Toxoplasma specific, 79
T-cell epitope, mycobacterial:
H-2 effects of flanking residues on recognition, 183
T-cell epitopes:
of house dust mite allergen Der p 2, 176
T-cell responses, regulation:
by IgE and IgG antibodies, 343
T-helper cell phenotypes:
in leprosy, 97
T-helper cell subsets:
human immunodeficiency virus replication, 85
Tolerogenic T-cell epitope:
suppression of arthritis, 110
Transforming growth factor-β:
down-regulation of major histocompatibility complex class I and increased susceptibility to natural killer cytokines, 263
Tuberculosis:
protection against by bone marrow cells expressing mycobacterial hsp65, 519
Tumour necrosis factor and interleukin-6:
modulation of oestrogen agonists and antagonists, 18
Tumour necrosis factor synthesis:
inhibition by chlorpromazine, 416
Tumour necrosis factor:
role in acquired resistance to *Listeria monocytogenes*, 256

V-region genes, of anti-Sm and anti-cardiolipin autoantibodies, 487